

ISSN : 2309-3234

DOI : 10.62948

TMSS Medical College Journal (TMCJ)

Volume 25, Issue 01, January 2026

Contents

Editorial

- ♦ From Immunohistochemistry to Molecular Testing for the Precision Cancer Management of Breast Cancer. 07-09
Rahman DA, Akter S.

Original Article

- ♦ Immunohistochemistry-Based Molecular Subtype Distribution of Invasive Breast Cancer and Its Clinicopathological Correlates in a Bangladeshi Cohort. 11-23
Rahman DA, Habib AA, Mazid A, Akter S, Ahmed F, Benzir M, Sakib MN, Rahman MT, Alamgir MH.
- ♦ Students' Perception of the Educational Environment Using the Dundee Ready Education Environment Measure (DREEM) at TMSS Medical College, Bogura, Bangladesh. 25-36
Habib MA, Hoque SMM, Ansari AAM, Tasnim T, Arzuman H.
- ♦ Menstrual Disorders and Relationship with Psychological Stress among Medical Students 37-43
Zohura FT, Rahman F, Jahan R, Alam MM, Ferdous R.
- ♦ Prognostic Significance of COL10A1 Overexpression in Breast Cancer Patients from Different Sociodemographic Backgrounds. 44-51
Roy NR, Das D, Hoque F, Jannat T, Tasnim A.
- ♦ Resource-Optimized PRP: A Comparative Study of Anticoagulants and Centrifugation Techniques in Bangladesh. 53-60
Hoque MA, Masud MAA, Basak SK, Mamun AA.
- ♦ Clinical Presentation and Culture Yield of Urinary Tract Infection Among Symptomatic Pregnant Women: A Tertiary Hospital Experience. 61-68
Sultana T, Akhther MH, Mahamuduzzaman AS, Shireen F, Parvin MR.



An Official Publication of TMSS Medical College



TMSS Medical College Journal (TMCJ)

January 2026

An Official Publication
of
TMSS Medical College, Bogura, Bangladesh

TMSS Medical College Journal (TMCJ)

Volume 25, Issue. 01, January 2026

An Official Publication of TMSS Medical College, Bogura, Bangladesh

Advisory Board

Prof. Dr. Hosne-Ara Begum
 Prof. Dr. Jawadul Haque
 Prof. Dr. Sayeba Akhter
 Prof. Dr. M. Afzal Hossain
 Prof. Dr. M. Aminul Islam
 Prof. Dr. Shamima Parvin Lasker

Editorial Board

Chairperson : Prof. Dr. Md. Zakir Hossain
Editor-in-Chief : Prof. Dr. Md. Shahjahan Ali Sarker
Associate Editor : Prof. Dr. Md. Abdul Mazid
Advisory Editor : Prof. Dr. Moudud Hossain Alamgir
Managing Editor : Dr. Md. Matiur Rahman

Local Members

Prof. Dr. A.K.M Ahsan Habib
 Prof. Dr. Md. Azfarul Habib
 Prof. Dr. Md. Rafiqul Islam
 Prof. Dr. Md. Anup Rahman Choudhury
 Prof. Dr. Nur-A-Farhana Islam
 Prof. Dr. Nahida Akther Zahan
 Prof. Dr. Tania Yeasmin
 Prof. Dr. Sultana Razia
 Prof. Dr. Shamim Ahmed Choudhury
 Prof. Dr. Md. Amirul Islam
 Dr. D. M Arifur Rahman
 Dr. Fahim Uddin Ahmad
 Dr. Md. Ahamedul Kabir

International Members

Prof. Dr. Dhananjaya Sharma
 Prof. Dr. Anurag Srivastava
 Prof. Dr. Shariq Rashid Mosoodi
 Prof. Olav Muurlink
 Prof. Dr. Paul Mailwaring
 Prof. Dr. Md. Faizus Sazzad
 Prof. Soraj Hongladarom
 Prof. Ravichandran Moorthy
 Prof. Darryl Macer
 Prof. Dr. Hasan Erbay

The TMSS Medical College Journal (TMCJ) is published two times a year (January and July). The TMSS Medical College Journal (TMCJ) is approved by Bangladesh Medical & Dental Council (BM&DC/4-D-2016/1478: Dated 15.01.2017). It accepts original articles, review articles and case reports. While every effort is always made by the Editorial Board and the members of the Journal Committee to avoid inaccurate or misleading information appearing in the TMSS Medical College Journal, information within the individual article are the responsibility of its author(s). The TMSS Medical College Journal, its Editorial Board and Journal Committee accept no liability whatsoever for the consequences of any such inaccurate and misleading information, opinion or statement.

Address of Correspondence

Editor-in-Chief, TMSS Medical College Journal, TMSS Medical College
 Rangpur Road, Thengamara, Bogura, Bangladesh. Phone: 01775-316617
 e-mail: editortmcjournal@gmail.com, Web: www.tmcjournal.org, www.tmssmedicalcollege.com

Printed by: TMSS Printing Press, Bogura, Bangladesh. e-mail: tmssprintingpress@gmail.com

TMSS Medical College Journal (TMCJ)

INFORMATION FOR AUTHORS

TMSS Medical College Journal (TMCJ) follows the guidelines and principles of the International Committee of Medical Journal Editors- ICMJE (www.icmje.org) and the COPE (www.publicationethics.org). Papers are accepted for publication with an understanding that they are submitted solely to the journal of 'TMSS Medical College Journal' and are subject to peer review and editorial revision. The statement, views and opinions expressed in the articles are those of the author(s) and are not necessarily those of the editor or publisher.

General Instructions

- a. Article must be within 5 years of research work.
- b. Authors of research work should be of relevant fields.
- c. Manuscript must be in English, Use Times New Roman 11 sized font.
- d. Type manuscripts double-spaced, on a good quality A4 sized paper, including references, figures with legends and tables on one side of the page only.
- e. Leave a margin of 05 cm on all sides of the page. Beginning with abstract page, all the pages (including pages containing tables, figures and references) will contain page numbers at the lower right-hand corner.
- f. Cite each reference in text in numerical order and list in the reference section.
- g. Begin each component of the manuscript with a new page in the sequence of (i) Title page with author information, name of the department(s) and institutions to which the work should be attributed, name and address of the author with post code responsible for correspondence and requests for reprints (ii) Abstract page (iii) Text (iv) References (v) Tables (vi) Figures with legends, for illustrations.
- h. Present the text in the sequence of Introduction, Methods & Materials, Results and Discussion.
- i. Use SI units of measurements (for blood pressure use mm of Hg).
- j. Define Statistical methods in method section of the paper.
- k. Use standard abbreviations, the full forms of which should precede in its first use in the text
- l. As a general rule, articles should not exceed 10-12 pages. Too much large volume manuscripts will not be accepted for publication.

Submission process

With two copies of manuscript (hard copy) directly or by courier or postal services to Editor-in-chief, TMSS Medical College Journal and/or A soft copy of Manuscript (MS Word) must be sent to editortmcjournal@gmail.com with scanned copy of all desired documents. Online submission will be possible through <https://www.tmcjournal.org>

Covering letter

Article must be accompanied by a covering letter, signed by all authors, containing statements that they are the only authors, The article represents original work and we take full responsibility for the information provided, the Manuscript has not been published elsewhere, accepted for publication elsewhere or under editorial review for publication elsewhere by any other journal. All copyright ownership of the manuscript will be transferred to the publishers of the TMSS Medical College Journal (TMCJ) once the article is accepted for publication in the journal. They must declare that they made the manuscript plagiarism free. On submission of the manuscript, the authors agree not to withdraw their manuscript at any stage prior to publication. Acknowledgements, Conflicts of interest & Funding are clearly mentioned, if any.

Author's contribution

In case of multiple authors, need an author contribution certificate (or statement) about specific contribution of each author in the study and the manuscript has been seen and approved as well as the order of authors in the manuscript.

Ethical Issues

Author must submit a copy of clearance certificate from an Institutional Ethical Review Board (IERB) or Institutional review Board (IRB) or Ethical Committee or from any authorized body along with the manuscript & mention the memo number within bracket in the methodology section while describing ethical issues.

Title page with author information (first page, but do not number)

- a. Full title of the article not exceeding 50 characters with three to five key words for use as indexing terms and as running title for use on top of the text pages.
- b. Author's names, highest academic degree, affiliations, and complete address for correspondence including Telephone/Mobile number, and e-mail address.

Abstract page (First numbered page)

- a. Do not cite references in the abstract. Be concise (250 words, maximum).
- b. Limit use of acronyms and abbreviations. Abbreviations must be defined at the first mention.
- c. The abstract should be a total of about 250 words maximum. Abstract should cover Background and Purpose (description of rationale for study); Methods (brief description of methods); Results (presentation of significant results); and Conclusion (succinct statement of data interpretation) in separate headings.

The Text/Main body

The following are typical main headings:

Introduction, Materials and Methods, Results, Discussion and Conclusion

- a. Introduction: Summarize the rationale for the study with pertinent references. The purpose(s) of the study should be clearly stated.
- b. Materials and Methods: Identify type of study and describe the study subjects and methods used. Provide methods of statistical analysis, Cite reference(s) for standard study and statistical methods. Describe new or modified methods. Identify apparatus (with name and address of manufacturer) used. Generic names of drugs must be given. Manuscripts that describe studies on humans must indicate that the study was approved by an Institutional Ethical Committee and that the subjects gave informed consent. Add statistical analysis.
- c. Results: Present only important results/observations in logical sequence in the text, tables or illustrations with relevant statistics.
- d. Discussion: Emphasize new and important results and the conclusions that follow including implications and limitations. Relate observations to other relevant studies.
- e. References: References must be listed following the Vancouver style. These should be numbered consecutively in the order in which they are first mentioned in the text as superscript number. Number of references should be kept below 40 except review article. All authors must be listed if less than six. Use et al, if more.
- f. Table & Figures: Tables & figures should be placed within the text of the manuscript. Number of tables & figures consecutively in the order of their first citation in the text. Tables should not duplicate in the text of the article and avoid internal horizontal & vertical rules. Total number of tables & figures must not exceed 10.

Original Article

Menstrual Disorders and Relationship with Psychological Stress among Medical Students

Zohura FT^{1*}, Rahman F², Jahan R³, Alam MM⁴, Ferdous R⁵

1. Dr. Fatema Tuz Zohura, Assistant Professor, Department of Physiology, TMSS Medical College, Bogura.
2. Dr. Farhana Rahman, Assistant Professor, Department of Physiology, Army Medical College, Jashore.
3. Dr. Rubaiya Jahan, Assistant Professor, Department of Physiology, Institute of Health Technology, Dhaka.
4. Dr. Md. Mashhurul Alam, Medical Officer, Department of Anesthesiology, Rajshahi Medical College, Rajshahi.
5. Dr. Rumana Ferdous, Assistant Professor, Department of Physiology, Rajshahi Medical College, Rajshahi.

Corresponding Author*Abstract**

Background: Menstruation is an inevitable part of a female's life and an important indicator of normal sexual and reproductive health. The changes in the normal menstrual pattern of female in the reproductive age may affect physical, physiological and psychological wellbeing. Stress may affect the pattern of menstrual cycle by stimulating the hypothalamo-pituitary-adrenal axis which is responsible for the production of cortisol hormone.

Objectives: The aim of the study was to evaluate menstrual disorders and their relationship with psychological stress among the students of Rajshahi Medical College, Rajshahi.

Materials and Methods: This cross-sectional type of descriptive study was carried out from January 2023 to December 2023 in the Department of Physiology, Rajshahi Medical College, Rajshahi. The study was conducted on 247 female students who were studying in Rajshahi Medical College after obtaining ethical clearance from the Ethical Review Committee of Rajshahi Medical College and informed written consent from the students. Data were collected by semi-structured questionnaire and analyzed by SPSS software, version-24. A p value < 0.05 was considered statistically significant for all tests.

Results: The students were 21.89 ± 1.44 years old on average, 12.01 ± 1.37 years old when they menarched, BMI 22.07 ± 3.34 kg/m², they had a monthly family income of 41874.49 ± 20436.26 taka, their menstrual cycle lasted 30.66 ± 6.43 days, and their flow lasted 5.26 ± 1.59 days. Marital status of the students showed that majority (89.50%) of the students were unmarried, (75.70%) of the students were from urban area and majority (85.40%) of the students lived in hostel. Of the students, 20.24% experienced oligomenorrhea, 6.88% hypomenorrhea, 7.29% menorrhagia, and 5.26% polymenorrhea in relation to menstrual disorders. The majority of students (77.30%) experienced moderate stress, followed by high stress (19.00%) and low stress (3.60%). Medical students' psychological stress levels were statistically significantly correlated with both menorrhagia and oligomenorrhea ($p < 0.01$ and $p < 0.001$, respectively). However, neither polymenorrhea nor hypomenorrhea was substantially associated with the medical students' psychological stress levels ($p > 0.05$ in both cases).

Conclusion: Psychological stress was significantly associated with the prevalence of various types of menstrual disorders. Timely diagnosis and management of menstrual abnormalities with proper counselling are necessary for improvement of health, sense of well-being, overall quality of life of students and lower the risks for future diseases.

Keywords: Medical students, Menstrual disorders, Premenstrual syndrome, Perceived stress scale and psychological stress.

Introduction

The normal reproductive years of females are characterized by monthly rhythmic changes in the rate of secretion of female hormones and corresponding physical changes in the ovaries and other sexual organs which is called menstrual cycle. A woman's menstrual cycle usually follows an average 28 day cycle.¹ A

normal menstrual cycle leads to formation of a dominant follicle with a mature ovum in the ovary then ovulation. In absence of fertilization, the endometrial lining sheds out resulting in discharge of blood from the uterus through the vagina which is called menstruation - a natural physiological phenomenon

occurring at a more or less regular monthly interval during women's reproductive life.²ovulatory, and luteal Menstruation first occurs in adolescents between 11 and 14 years of age with a period length of average 7 days with a cycle length ranging from 21 to 45 days with an average blood loss of 20 to 80 ml.³ This phenomenon is not an easy one and is often associated with some degree of suffering. According to several studies, a large proportion of the female population of reproductive age suffer from menstruation related health issues. There are various forms of menstrual disorders. The list of menstrual disorders ranges from dysmenorrhea, premenstrual syndrome (PMS), polymenorrhea, oligomenorrhea, menorrhagia, amenorrhea to irregular menstrual cycle. Dysmenorrhea is the commonest gynecological disorder among women with prevalence of 60% to 93%.⁴

Other abnormalities are also common. Although menstrual disorders are common health problems, they are generally perceived as only minor health concerns and thus irrelevant to the public health agenda particularly for females in developing countries.⁵ But menstrual abnormalities not only cause deterioration in quality of life but also have detrimental effects on mental and reproductive health and also carry economic burden. Menstrual disorders over prolonged periods of time can cause anovulation, endometrial hyperplasia, infertility,⁴ osteoporosis, breast cancer.⁶ Menstrual patterns can be affected by various factors including age, ethnicity, family history, physical activity, dietary habits. Stress can be a major contributor to or cause of menstrual irregularities.⁷

Several studies have shown association of stress with variation in the length of menstrual cycle, anovulation, duration and amount of menstrual bleeding. Multiple pathways have been proposed linking perceived stress with reproductive function. The hypothalamic pituitary adrenal axis is activated during stress that suppresses hypothalamic gonadotropin releasing hormone (GnRH) secretion. GnRH stimulates the production of follicle stimulating hormone (FSH) and luteinizing hormone (LH) which activate the ovary to secrete estradiol and progesterone. Suppression of GnRH

secretion suppresses reproductive hormone secretion and also leads to a delay or inhibition of the LH surge. Thus stress can alter hormonal profiles and in turn menstrual outcomes.^{6,8}

Stress is prevalent among medical professionals and that is also true for the medical students who have very busy and demanding schedules. Psychological stress is common among medical students for various reasons. High expectations of parents, exhausting work hours, continuous evaluation process, striving for earning high grades, enormous syllabus to be covered in a limited time period, thought of appearing or failing in exams, performance in periodic examinations are potential sources of stress among medical students.⁹ Due to the high prevalence of psychological stress among them, medical students are at high risk of developing menstrual abnormalities. Several studies have been done in different countries regarding the proportion of menstrual disorders and their association with psychological stress among the medical students but to the best of our knowledge very few studies has been done in our country. Therefore, the study was designed to assess the proportion of medical students suffering from menstrual disorders and their relationship with psychological stress in our context.

Materials and Methods

This study was a cross-sectional descriptive study. It was conducted in the Department of Physiology, Rajshahi Medical College, Rajshahi during the period of January 2023 to December 2023. Female students of Rajshahi Medical College were enrolled in this study. Before the starting of the study, a questionnaire was prepared according to the objectives of the study. All questions regarding the participants demographics as well as their menstrual pattern (menarche age, flow duration in days and its quantity, cycle length in days) and history of oligomenorrhea, polymenorrhea, hypomenorrhea and menorrhagia were included in the questionnaire. For assessment of stress level, Perceived Stress Scale was used. Then on the basis of inclusion and exclusion criteria history was taken from the students. Individuals who were match according to the selection criteria of the study were informed about

the purpose of the study. After being informed about the purpose of the study individuals who gave consent to participate in the study were finally selected as a study subject. Then data were collected based on variables of interest.

Sampling:

Stratified random sampling technique was applied to select each individual in this study. As we knew that in Rajshahi Medical College, total 4 phases medical students are available. In this study, students were divided into 4 strata-131 female students in phase 1, 122 students in phase 2, 109 students in phase 3 and 134 students in phase 4. So, total population was 496. As sample size was 247, sampling fraction = $[(247 \div 496) \times 100] = 49.80\%$. So, 65 students from phase 1, 61 students from phase 2, 54 students from phase 3 and 67 students from phase 4 were selected.

Inclusion Criteria:

Female students of Rajshahi Medical College, Rajshahi had been studying medical science for at least 1 year.

Exclusion Criteria:

Medical students with known chronic health problems such as thyroid disorders, students with any history of psychiatric problems, diagnosed pelvic pathology such

as fibroids, endometrioses, polycystic ovarian syndrome, adenomyosis, any pregnant and lactating mother among the medical students, hormonal contraceptives users were excluded from the study.

Data Collection Tools:

The data were taken with the help of using a semi-structured questionnaire (research instrument) containing all the variables of interest. For assessment of psychological stress level Perceived Stress Scale was used.

Statistical Analysis:

All data were analyzed by using the 'Statistical Package for Social Sciences (SPSS)' software, 24-version. A Chi-square test was used to observe the relationship of menstrual disorders with the level of psychological stress among the students. A p-value < 0.05 was considered statistically significant for all tests.

Institutional Review Board Statement:

Permission was taken from the Institutional Review Board (IRB) of Rajshahi Medical College (RMC), Rajshahi before the starting of the study regarding the study design, methodology and target population characteristics.

Results

Table I: Distribution of the students by their different variables (n=247).

Variables	Mean $\pm$ SD	Range
Age (Years)	21.89 $\pm$ 1.44	19 – 25
Age of menarche (Years)	12.01 $\pm$ 1.37	8 – 16
BMI (kg/m ²)	22.07 $\pm$ 3.34	14.12 – 34.35
Monthly family income (taka)	41874.49 $\pm$ 20436.26	10,000-1,00,000
Menstrual cycle length (days)	30.66 $\pm$ 6.43	15-48
Duration of flow (days)	5.26 $\pm$ 1.59	1-11

Table I shows the mean age of the students was 21.89 $\pm$ 1.44 years, age of menarche was 12.01 $\pm$ 1.37 years, BMI 22.07 $\pm$ 3.34 kg/m², monthly family income 41874.49 $\pm$ 20436.26-taka, menstrual cycle length 30.66 $\pm$ 6.43 days and duration of flow 5.26 $\pm$ 1.59 days.

Table II: Marital status, residence and present living situation of the student (n=247).

Variables	Frequency	Percentage (%)
Marital status		
Married	26	10.50
Unmarried	221	89.50
Residence		
Rural	60	24.30
Urban	187	75.70
Present living situation		
With family	36	14.60
Hostel	211	85.40

Table II reveals that 89.50% of students were single and 10.50% married. Most students (75.70%) were urban and 24.30% rural. Currently, 85.40% of students live in hostels and 14.60% with families.

Table III: Distribution of the students by presence of various menstrual disorders and psychological stress level (n=247).

Menstrual disorders	Frequency		Percentage (%)
Oligomenorrhea	Yes	50	20.24
	No	197	79.76
Polymenorrhea	Yes	13	5.26
	No	234	94.74
Hypomenorrhea	Yes	17	6.88
	No	230	93.12
Menorrhagia	Yes	18	7.29
	No	229	92.71
Levelsofpsychological stress	Low stress	9	3.60
	Moderate stress	191	77.30
	High stress	47	19.00

Table III demonstrates that 20.24% of students had oligomenorrhea, 6.88% hypomenorrhea, 7.29% menorrhagia, and 5.26% polymenorrhea. Most of the students (77.30%) were moderately stressed. 19.00% had high and 3.60% had low stress.

Table IV: Relationship between the menstrual disorders and level of psychological stress among the students (n=247).

Variables	Levels of psychological stress			p-value
	Low stress	Moderate stress	High stress	
Oligomenorrhea	1 (11.10%)	25 (13.10%)	24 (51.10%)	< 0.001
Polymenorrhea	0 (0%)	10 (5.20%)	3 (6.40%)	> 0.05
Hypomenorrhea	1 (11.10%)	12 (6.30%)	4 (8.50%)	> 0.05
Menorrhagia	0 (0%)	9 (4.70%)	9 (19.10%)	< 0.01

In Table IV, oligomenorrhoea and menorrhagia were found to significantly correlate with psychological stress in medical students ($p < 0.001$ and $p < 0.01$, respectively). Both hypomenorrhea and polymenorrhea did not significantly affect psychological stress in medical students ($p > 0.05$).

Discussion

The average age of students in this study was 21.89 ± 1.44 years. Alshdaifat et al.¹⁰ observed similar results, with a mean student age of 21.6 ± 2.2 years. Nagma et al.¹¹, Alshayeb and Sakka¹², and Khaleel¹³ observed similar results. Kollipaka et al.¹⁴ obtained different results.

This study found that 89.50% of students were single and 10.50% were married. Alshayeb and Sakka¹² reported 87.0% of students were single. Alshdaifat et al.¹⁰, Alhammadi et al.¹⁵, and Rafique and Al-Sheikh⁷ reported comparable results. In this study, 85.40% of students lived in hostels and 14.60% with families. Alshdaifat et al.¹⁰ and Alshayeb and Sakka¹² found different results.

In the study, students had a mean BMI of 22.07 ± 3.34 kg/m². Khaleel¹³ reported similar results, with a mean BMI of 22.71 ± 6.369 kg/m². The mean student BMI in Nagma et al.¹¹ was 20.91 kg/m², which contradicted the findings. Different eating habits in different regions may explain the differences.

The mean age of menarche in this study was 12.01 ± 1.37 years. Cakir et al.¹⁶ observed similar results, with a mean menstrual age of 12.8 ± 1.3 years. Singh et al.¹⁷, J et al.¹⁸, Alshdaifat et al.¹⁰, Alhammadi et al.¹⁵, Khaleel¹³, and Rigon¹⁹ found similar results. In contrast, Esimai and Esan⁵ observed a mean menarche age of 14.2 years. Mishra and Mukhopadhyay²⁰ and Verma et al.²¹ observed conflicting results.

This study found a mean menstrual blood flow length of 5.26 ± 1.59 days. Similarly, Lakkawar et al.⁴ found a mean menstrual blood flow length of 4.25 ± 0.5 days. Climate and seasonal change may explain these differences. Adolescent females suffer more from

hormone imbalances and longer or more frequent periods in summer. The study found 20.24% of students had oligomenorrhea. Lakkawar et al.⁴ observed 21% of students experienced oligomenorrhoea, like the study. Alshayeb and Sakka¹² discovered oligomenorrhea in 5% of students, lower than the study. Research found 7.29% of students experienced menorrhagia. Alshayeb and Sakka¹² observed 7% of students experienced menorrhagia, like the study. Alshayeb and Sakka²² found 8.6% of students had menorrhagia, greater than the research. Also, Houston et al.²³ obtained different results.

The recent study found 5.26% of medical students had polymenorrhea. Lakkawar et al.⁴ discovered that 8% of students had polymenorrhoea. Rafique and Al-Sheikh⁷ discovered polymenorrhoea in 13% of pupils.

The study found that 77.30% of students had moderate stress, 19.00% had high stress, and 3.60% had low stress, with a mean perceived stress scale score of 22.03 ± 4.78 . Khaleel¹³ obtained similar results. In contrast, Achmad et al.²⁴ observed that 55.8% of students had severe stress, 36.5% moderate, 5.8% extremely severe, and 1.9% mild. Kollipaka et al.¹⁴ obtained different results.

A strong correlation ($p < 0.001$) was identified between oligomenorrhoea and psychological stress among students in this study. Alshayeb and Sakka¹² obtained similar results.

This study observed no correlation between hypomenorrhea and student psychological stress ($p > 0.05$). Nagma et al.¹¹ observed similar results. Researchers found a substantial correlation between menorrhagia and psychological stress in medical students ($p < 0.01$). Khaleel¹³ observed a statistically significant association between menorrhagia and psychological stress in medical students. Khaleel¹³ obtained similar results. However, Zhou et al.²⁵, Sood et al.²⁶, Nagma et al.¹¹, and Fenster et al.⁶ observed different results. Different study methodology and regional variance may explain these inconsistencies.

This study revealed no correlation between polymenorrhea and student psychological stress ($p>0.05$). Alshayeb and Sakka¹² obtained similar results.

Conclusion

There was a really high prevalence of stress among these students and it was also associated with the occurrence of various menstrual disorders. So, all medical faculties should set up procedures for the early detection of students with stress and menstruation disorders which might be helpful for prevention of various diseases.

Funding

The authors received no financial support for the research, authorship and/or publication of this article.

Conflict of interest

Authors declared no conflict of interest.

References

1. Hall JE, Hall ME. Guyton and Hall textbook of medical physiology; Second South Asian Edition. India: Elsevier; 2015.
2. Itriyeva K. The normal menstrual cycle. *Curr Probl Pediatr Adolesc Health Care*. 2022 May 6;52(5):101183.
3. Periasamy P, Suganthi V, Karuppiah P. Association of menstrual patterns with perceived stress score in college-going female students of a South Indian Town. *Apollo Medicine*. 2021; 8 (2): 76-79.
4. Lakkawar N, L. J, Arthi N, Alagan P, N V. A Study of Menstrual Disorders in Medical Students and its Correlation with Biological Variables. *Scholars Journal of Applied Medical Sciences*. 2014; 2 (6E): 3165-3175.
5. Esimai O, Esan GO. Awareness of menstrual abnormality amongst college students in urban area of ile-ife, osun state, Nigeria. *Indian J Community Med*. 2010 Jan;35(1):63–66.
6. Fenster L, Waller K, Chen J, Hubbard AE, Windham GC, Elkin E, et al. Psychological stress in the workplace and menstrual function. *Am J Epidemiol*. 1999 Jan 15;149(2):127–134.
7. Rafique N, Al-Sheikh MH. Prevalence of menstrual problems and their association with psychological stress in young female students studying health sciences. *Saudi Med J*. 2018 Jan;39(1):67–73.
8. Schliep KC, Mumford SL, Silver RM, Wilcox B, Radin RG, Perkins NJ, et al. Preconception Perceived Stress Is Associated with Reproductive Hormone Levels and Longer Time to Pregnancy. *Epidemiology*. 2019 Nov;30 Suppl 2:S76–S84.
9. Ray I, Joseph D. Stress in medical students. *JK Science*. 2010; 16:47.
10. Alshdaifat E, Absy N, Sindiani A, AlOsta N, Hijazi H, Amarin Z, et al. Premenstrual Syndrome and Its Association with Perceived Stress: The Experience of Medical Students in Jordan. *Int J Womens Health*. 2022 Jun 14;14:777–785.
11. Nagma S, Kapoor G, Bharti R, Batra A, Batra A, Aggarwal A, et al. To evaluate the effect of perceived stress on menstrual function. *J Clin Diagn Res*. 2015 Mar 1;9(3):QC01–3.
12. Alshayeb S, Sakka K. The prevalence of various menstrual disorders and its association with psychological stress in medical faculties students. A Cross-sectional study. *Res Sq*. 2022 Dec 9; 20 (2): 10-15.
13. Khaleel, D. The Effect of Psychological Stress on Menstrual Cycle among Medical Students. Ministry of Higher Education and Scientific Research, 2021; 6 (1): 1-17.
14. Kollipaka R, Arounassalame B, Lakshminarayanan S. Does psychosocial stress influence menstrual abnormalities in medical students? *J Obstet Gynaecol*. 2013 Jul;33 (5): 489–493.

15. Alhammadi M, Albogmi A, Alzahrani M, Shalabi B, Fatta F, AlBasri S. Menstrual cycle irregularity and examination stress among female medical students at king abdulaziz university, jeddah, saudi arabia. *Res Sq.* 2022 Feb 23;
16. Cakir M, Mungan I, Karakas T, Giriskan I, Okten A. Menstrual pattern and common menstrual disorders among university students in Turkey. *Pediatr Int.* 2007 Dec;49(6):938–942.
17. Singh A, Kiran D, Singh H, Nel B, Singh P, Tiwari P. Prevalence and severity of dysmenorrhea: a problem related to menstruation, among first and second year female medical students. *Indian J Physiol Pharmacol.* 2008 Dec;52(4):389–397.
18. J B, AM H, SA N. Menstrual Pattern and Common Menstrual Disorders among Students in Dinajpur Medical College. *Dinajpur Med Col J.* 2009; 2 (2): 37 43.
19. Rigon F, De Sanctis V, Bernasconi S, Bianchin L, Bona G, Bozzola M, et al. Menstrual pattern and menstrual disorders among adolescents: an update of the Italian data. *Ital J Pediatr.* 2012 Aug 14;38:38.
20. Mishra SK, Mukhopadhyay S. Socioeconomic correlates of reproductive morbidity among adolescent girls in Sikkim, India. *Asia Pac J Public Health.* 2012 Jan;24(1):136–150.
21. Verma D, Pandya D, Ramanuj D, Singh D. Menstrual Pattern of Adolescent School Girls of Bhavnagar (Gujarat). *NJIRM.* 2011; 2 (1): 38-40.
22. Karout N, Hawai SM, Altuwaijri S. Prevalence and pattern of menstrual disorders among Lebanese nursing students. *East Mediterr Health J.* 2012 Apr;18(4):346–352.
23. Houston AM, Abraham A, Huang Z, D'Angelo LJ. Knowledge, attitudes, and consequences of menstrual health in urban adolescent females. *J Pediatr Adolesc Gynecol.* 2006 Aug;19(4): 271-275.
24. Achmad L, Sirait B, Semen G. The relationship between stress levels and menstrual cycle regularity in students of the faculty of medicine, universitas kristen Indonesia, class of 2017. 7 (8): 74-79.
25. Zhou M, Wege N, Gu H, Shang L, Li J. Work and Family Stress is Associated with Menstrual Disorders but not with Fibrocystic Changes: Cross-sectional Findings in Chinese Working Women of occupational health. 2010; 52 (6): 361-366.
26. Sood M, Devi A, Mohd Daher AA, Razali S, Nawawi H, Hashim S, et al. Poor Correlation of Stress Levels and Menstrual Patterns among Medical Students. *jABs.* 2017 Nov 4;2(5):73–78.